


Studies of the ECR plasma in the visible light range

S. Biri , R. Rácz, J. Pálinkás*



Institute of Nuclear Research (ATOMKI), Debrecen, Hungary

** University of Debrecen, Dept. Exp. Physics, Hungary*



Outline

- Motivation
 - Experimental setup
1. ECRIS settings effects
 2. Cold and warm electrons
 3. The color of plasmas (Xe, He)
- Conclusion

Motivation

ECR plasmas can be investigated by different ways



Electrostatic (Langmuir) probes


- Normal, double, emission types etc.
- Local information
- Plasma density, potential
- Technical difficulties

EM-radiation recording

- Ranges: IR, VL, UV, XR
- Imaging techniques (photos)
- Spectrum analysis
- Integrated axial information

Computer simulations

- Different methods
- Many nice graphical results
- Initial conditions...
- Prove: comparison with experiment



XR-region: ion excitation, brehmstrahlung:
plasma ions and lost electrons
VL-region: atom excitation: cold plasma electrons

VL-region: Why? (an ECRIS is not a light source...)

There are at least two reasons worth studying the VL-part of ECR plasmas.

1.

- The cold electron population is the “first product” of the step-by-step ionization process.
- To get warm and hot electrons **later** we need cold electrons **first**.
- Cold electrons: starting phase toward the high charge ionization.


2.

- The application area of the ECR ion sources is broadening. Main field is still to produce highly charged ions (**HCI**).
- ECR-heating principle (+ B-minimum): **LCI** high current beams (e.g. proton, carbon) also by ECR ion sources.
- **Medical** applications, European Spallation Source (**ESS, proton**).
- In such sources the energy of the electron population is much lower than in the traditional **HCI** ECRISs.

Therefore we made high resolution ECR plasma photo series and movies


- Motivation
- **Experimental setup**
- ECRIS settings effects
- Cold and warm electrons
- The color of plasmas (Xe, He)
- Conclusion

The ATOMKI-ECRIS




PLASMA CHAMBER

EXTRACTION



Experimental setup



Layout of the experiment, top view. Plasma chamber side view (1), plasma chamber end wall (2), resonance zones (3, 4), waveguides (5,6), bias disc (7), gas tube (8), mirror (9), camera (10). At bottom the axial magnetic field distribution made by middle-powered solenoids, is shown (11).


Camera settings

- The ECR plasma is not an ideal photo model.
- Its longitudinal length is about 20 cm.
- It is partly transparent and diffuse.

- Cameras: Canon A630 and Sony HDR-FX7E
- Picture size: 8 MP
- Exposure time: 0.8-4 sec
- Iris value: 8
- ISO value: 80
- Distance: $100+40=140$ cm




What we see...




**Plasma „spider“: only 6 legs, not 8
The 3+3 legs or arms fed by bunches of loss lines.**


Operation modes



+



=




*14.3 GHz, Klystron,
10-100 W*

*8.4-12.4 GHz
TWT, 5-20 W*

*Klystron+TWT,
double plasma*

- Motivation
- Experimental setup
- **ECRIS settings effects**
- Cold and warm electrons
- The color of plasmas (Xe, He)
- Conclusion


ICIS09 poster, R. Rácz et al., (RSI 81, 2010, 02B708)



TrapCAD simulation

Effect of the microwave power



Neon



Argon

Gas dosing rate

Neon(mbar) 8,5E-6 7,0E-6 2,5E-6 1,1E-6 8,0E-7 5,0E-7 4,4E-7



Residual gas plasma
(valves closed).

15 sec exp. time

*YouTube, ANL ECRIS movie, 8 min.
More than 3000 visitors since 2007!*

- Motivation
- Experimental setup
- ECRIS settings effects
- **Cold and warm electrons**
- The color of plasmas (Xe, He)
- Conclusion

Cold and warm electrons study

(some results submitted to Plasma Sources Science and Technology)

ECR plasma electron components: 3 **artificial** populations

cold electrons:
1-200 eV

warm electrons:
several keV

hot electrons:
hundreds of keVs

Comparison:

Visible-light photo

AND

Simulation

Comparison:


X-ray photo

AND

Simulation

(do not deal here)

Comparison: VL-photo and electron simulation




- $5 \cdot 10^5$ starting particles
- start from resonance zone
- start energy: 1-100 eV
- random direction
- ECR heating ON
- Calculation time: enough statistics for lost and non-lost electrons
- filtered energy: 1-200 eV
- 5000 electrons left

*Typical argon plasma.
Photons from excited atoms
and ions. 14 GHz, 50 W*

*TrapCAD simulation,
14 GHz, electrons*

- **Good agreement** between the photons and electrons.
- Cold electrons are not so well bounded: the plasma is **not empty**.
- In contrast to high energy electrons, cold electrons are very effective in **exciting** visible light.
- VL-photographs show the plasma region with cold electrons **confined almost equally** inside the RZ.

Comparison: electron simulation and X-ray-photo



- Good agreement between simulation and XR-photo.

- Warm electrons are trapped at magnetic **gap**.


- Argon ions locate at the **same** positions.

- Strong azimuthal and radial inhomogeneity.


TrapCAD simulation, 14 GHz, warm electrons. The same output file was used as for cold electrons. Filtering here: 3-10 keV

X-ray photo, argon $K\alpha$ radiation (cca 3 keV) 14 GHz, 50 W

S. Biri, ECRIS2010, Grenoble





Comparison of density profiles



Particle density profiles from the center toward the magnetic *pole and gap*.

Simulation: 3-10 keV warm electrons

Measurement: 3 keV X-ray photons from argon ions



- Calculation and photo clearly show strong azimuthal and radial inhomogeneity of the plasma.
- The central plasma density is much lower for the case of ions and warm electrons than for cold electrons and VL-photons.
- This information is very important for future ion source designers (extraction optical system).

- Motivation
- Experimental setup
- ECRIS settings effects
- Cold and warm electrons
- **The color of plasmas (Xe, He)**
- Conclusion

The color of the ECR plasmas

It is a challenging task to **understand** the color of different ECR gas plasmas.


The color of the plasma can be determined by visible light electron **transitions** of plasma components (atoms and ions).







Examples: **Xe** spectrum (CRC Handbook of Chemistry and Physics 2009).

Electromagnetic radiation in the VL-range from 360 nm (violet) to 820 nm (red)


Human eye peak sensitivity: the spectral luminous efficiency function (SLEF) peaks at 555 nm (green)

Xenon spectrum normalized with SLEF.







- There is a good **visual** agreement between the calculated normalised color and the real color of the plasmas.
- Also there is good agreement between the **RGB values** of the decomposed normalized spectrum lines and of the photos.
- Thus this process is able to **explain** and **understand** the color of ECR plasmas.




Krypton, Xenon and mixed (50%) plasmas.



Neon, Argon and mixed (50%) plasmas.




Nitrogen, Oxygen and mixed (50%) plasmas.



Methane, Helium and mixed (50%) plasmas.



Conclusion

- Visible light (VL) photos transform information mainly on the cold electron component of the plasma. Cold electrons are confined in the central plasma part.
 - X-ray (XR) photos show the spatial distribution of ions. These ions and the warm electrons are well confined by the magnetic field lines structure showing strong asymuthal and radial inhomogenity.
 - The color of the ECR plasmas can be explained and understood by the atomic transitions combined with human eye sensitivity.
 - We are convinced that VL and XR photos hide many more interesting and valuable information on the ECR plasma...
- 

Thank you for your attention!