



# Effect of self-consistency on periodic resonance crossing

G. Franchetti

HB 2012

Beijing, China

# Overview

Frozen approach to long term space charge simulations

Mechanism, simulations, benchmarking

Markovian update, Markovian mapping

Behavior at limit  $n \rightarrow \infty$


Summary / Outlook

# Simulation challenges

**Long term tracking** of **high intensity beams** enhances the code limitations

PIC simulations are affected by noise, which may compete with the physical mechanisms one tries to simulate


**Cure:** use large amount of macro-particles → **only short term tracking feasible**



**Work started with “frozen” space charge under the assumption of small beam loss**

Frozen tracking suitable in case of small beam loss prediction → **no noise effect!**  
*(we certainly do not want operate a machine in a regime of large beam loss!)*

# Beam loss mechanism: resonance crossing



# 成份 = ingredients

Effects caused by a periodic resonance crossing is determined by the following factors:

1. Distance of the bare tune from the resonance
2. Space charge tune-spread (main detuning source)
3. Speed of resonance crossing (synchrotron frequency)


Speed of resonance crossing is linked to the strength of the resonance:

Larger crossing speed through strong resonances make the similar effect of smaller crossing speed through weak resonances

# Effect of resonance crossing

- 1) Space charge stabilize resonant behavior
- 2) Particles follow a dynamics determined by fixed points (trapping)  
Trapping originally studied by A.Chao 1976



In the case of non-adiabatic crossing



If the attraction point exists, then the particles are carried away by the attraction point (trapping) otherwise they are **scattered** by the resonance crossing

# Experimental verification in SIS18 2008-2010

Code benchmarking, but also proof of principle experiment





G. Franchetti, O. Chorniy, I. Hofmann, W. Bayer, F. Becker, P. Forck, T. Giacomini,  
M. Kirk, T. Mohite, C. Omet, A. Parfenova, and P. Schuett PRSTAB **13**, 114203 (2010)

# Application to SIS100: frozen approach



Typical pattern of beam loss for a frozen beam tracking



Frozen model: means that the source of detuning with amplitude remains unaffected by the beam loss




# Correcting 2<sup>nd</sup>, and 3<sup>rd</sup> order, normal and skew resonances




# The issue of self-consistency

At the beginning of storage




# The issue of self-consistency

After some beam loss




# The issue of self-consistency

When the tune spread will not cross the resonance the beam loss process stops



# Consequences?

Potentially the most dangerous situation is when the tune is close to the resonance, because most of the particles composing the beam would be lost



# Markovian ansatz

**Ansatz:** we assume that the transverse particle distribution is of the same type any time and that the emittances does not change.  
However, the beam intensity is updated.

This represents a kind of *half* self-consistent approach → only the intensity is updated


The ansatz that **the distribution does not change** means that the particles are lost from everywhere inside the beam distribution....



## Idea

The Markovian update is at each step made by piece of a frozen space charge update

# Why Markovian? The loss of memory



# Simulations

## Simulation method:


For speeding up simulations several clones of the same simulations are run in many processors, and each simulation tracks 4 macro-particles (used in SIS100 studies)

**By applying a Markovian update,  
how many particles should we use ????**

**If in a simulation with 4 macro-particles one particle is lost, that corresponds to a change of 25% of the intensity and certainly the results cannot be meaningful....**

# Markovian update for SIS100

Taking a systematic approach we look at what happen by taking a case with 4, 10, 20, 100 macro-particles



# Markovian Mapping

Assume in the frozen simulations  
the beam intensity evolves as

$$\frac{dI}{dt} = -\Delta(I_0)f\left(\frac{t}{\tau(I_0)}\right) \frac{1}{\tau(I_0)}$$



This is a fit that can be always be  
made



The Markovian update takes  
the form

$$\frac{dI^*}{dt} = -\Delta(I^*)f\left(\frac{t(I^*)}{\tau(I^*)}\right) \frac{1}{\tau(I^*)}$$



Markovian mapping

# Example with SIS100 simulations

For example, by taking the ansatz that


$$I = \frac{I_0}{1 + [t/\tau(I_0)]^{1.23}}$$

we fit the curves of beam survival obtained from the “frozen” simulations



Rough approach but is a start for checking the concept

The function  $\tau(I_0)$  it is found by fitting to be  $\tau(I_0) = 2.5 I_0^{-3.3}$  with  $I_0$  in units of  $10^{11}$




Therefore integrating the equation for  $I^*$  we find

$$I^* = I_0 [1 + 0.87 I_0^{3.3} t]^{-1/3.3}$$

with  $t$  in units of 1000 turns  
and  $I_0$  in units of  $10^{11}$  atoms

# Comparison with Markovian update



← very good agreement!

$I_0 = 0.25 \times 10^{11}$

← Beam survival for Markovian update.  
In green the curve  $I^*$  from the  
Markovian process

← Beam survival with frozen  
space charge

quite promising approach !!!

# Advantages of the approach

For making a self-consistent simulations a large number of macro-particles should be used in order to “smooth” the beam loss process



But that would require long time unless a development of parallel computing


The discussed properties allows one can compute fast result from frozen space charge simulation for several intensities and later using a Markovian mapping find a result similar the full simulations

How does beam survival evolves for very long term simulations ?



We can study this problem by using Markovian-update simulations

# Intuitively expected behavior



# The asymptotic limit: case of a single resonance

DQx = 0.15  
turns  $\rightarrow 5E6$   
pipe at 4.5 sigma

The “close to the resonance collapse” does not happen


**Why ?**

**Close the resonance** the Markovian update simulation overlaps with the frozen simulation



**Is it self-consistency not important in some condition?**

Full simulation with a Markovian update



Difficult issue:  
**what is the time scale of the process ?**

# Summary / Outlook

Self consistency brings some interesting feature

- 1) Self-consistency was studied in a Markovian approach
- 2) In this approach self-consistency seems to mitigate the impact on beam losses on SIS100
- 3) An unexplained robustness of the Markovian update is found !
- 4) For a single resonance the “close to the resonance collapse” does not happen !!
- 5) Markovian mapping seems a promising tool to make predictions

## Outlook

- 1) Include in the method a re-update of rms sizes
- 2) Comparison with 2D5 PIC simulations (short scale)

Further benchmarking with experimental results

**New benchmarking with CERN-PS data.  
1<sup>st</sup> measurements performed in June 2012**



Thanks for the attention