



**24th International Workshop on ECR Ion Source
(ECRIS'20)**
28-30th, September, 2020

Numerical Simulations of Plasma Dynamics in ECRIS Afterglow

Li Lei¹, Xiaolin Jin^{1*}, Jibo Li², Tao Huang¹, Zhonghai
Yang¹ and Bin Li¹


1. *Vacuum Electronics National Laboratory, University of Electronic Science and Technology of China, Chengdu*
2. *Institute of Modern Physics, Chinese Academy of Sciences, Lanzhou*

Email address: jinxiaolin@uestc.edu.cn



ECRIS Set-up

Layout of LAPECR1U



Critical Parameters

Cavity Radius	19 mm
Cavity Length	78 mm
Aperture Radius	6 mm
MW Mode	$TE_{1,1}$
MW Frequency	14.5 GHz
MW Power	500 W
ECR Magnetic Field	0.518 T
Plasma Species	Argon
Plasma Density	10^{12} cm^{-3}



Simulation Scheme

Two-Step Simulation for A Stable Operation

Energetic Electrons

- Single Particle Simulation**
 - Electrons (~10 eV) uniformly distributed at first
 - Coulomb Collision is considered with constant scattering factor
 - Rejoin system as new particles when out of bounds
 - Electrons are heated till average energy reaches 10 keV
 - Distributions of the electrons are sent to the 2nd step as initiation

ECRIS Operation



- Implicit Particle-in-cell Simulation**
 - Cold Electrons (~10 eV) uniformly distributed at first
 - Ions (~1 eV) distributed due to quasi-neutrality
 - Coulomb Collision is considered with computed local plasma density
 - Secondary emission for electrons and absorption for ions when out of bounds
 - Ionization is omitted, and instead new particles are manually created





ECRIS Potential at CW Operation

*Energetic electrons (well-confined)
as backgrounds instead of being tracked*

**With
MW Power**



**Without
MW Power**




MW power is critical to the formation of the potential dip.




Cold Electrons at Power-off


assuming that the energetic electron distribution still remain unchanged



Density Distribution



Power Off





Cold electrons by itself may not contribute much to the ECRIS afterglow.




Energetic Electrons at Power-off

Single particle simulation



Density Distribution
↓
Power Off



Energetic electrons are less confined at power-on due to RF diffusion.



Summary

Next Works:

- I. Track energetic electrons in the simulation once an appropriate velocity distribution has been calculated.
- II. Consider multi charge state ions and step-wise ionization.
- III. Extend simulation time for diagnostics of ion dynamics.

Thanks for your attention!