

# Spin-Tracking Simulations in a Proton Prototype EDM Ring using Bmad

M. Vitz<sup>1</sup>, A. Lehrach<sup>1,2</sup>, Institut für Kernphysik 4, Forschungszentrum Jülich, 52425 Jülich, Germany,

R. Shankar, Istituto Nazionale di Fisica Nucleare, sezione di Ferrara, 44122 Ferrara, Italy


<sup>1</sup>also at III. Physikalisches Institut B, RWTH Aachen University <sup>2</sup>and JARA-FAME, 52056 Aachen, Germany

on behalf of the JEDI Collaboration

## Physical Motivation

- **Electric Dipole Moment (EDM)** is a fundamental property of a subatomic particle, similar to the Magnetic Dipole Moment (MDM).

- Source of  $\mathcal{P}$  and  $\mathcal{T}$  violation ( $\overset{CPT}{=} CP$  violation) and therefore closely connected to **matter antimatter asymmetry**.




- EDM of charged particles can be measured in a storage ring as spin rotation is defined by EDM and MDM contribution [1].

- **Vertical spin build-up** due to EDM needs coherent spin rotation for  $\sim 1000$ s in the horizontal plan due to MDM.

- Spin tracking simulations with Bmad Software Library [2] to estimate amount of time spins are aligned.  
⇒ **Spin Coherence Time (SCT)**

## Prototype Design


- The Proton Prototype EDM Ring (PTEDM) **combines E and B** fields so the ref. particle spin is aligned to the momentum [3].



- Particle Species: Proton
- Ref. Momentum: 294 MeV
- Circumference: 123.358 m
- Dipole E-Field: 5.061 MV/m
- Dipole B-Field: 0.023 T

- Design contains two active quadrupole families (QD and QF) with sextupoles (family SXQF and family SXQD) placed on them.

—  $\beta_x$  [m] —  $\beta_y$  [m] —  $D_x$  [m] —  $D_y$  [m]






- $Q_x = 1.823$
- $Q_y = 1.123$
- $\xi_x = -0.070$
- $\xi_y = +0.035$

- To estimate SCT a fully longitudinal polarized bunch consisting of  $10^3$  particles was tracked for  $10^6$  revolutions ( $\sim 1$ s) in the ring.

## Spin Tracking

- Spin depolarization can be optimized by adjusting system parameters like the **chromaticities  $\xi_x$  and  $\xi_y$** .
- Chromaticities can be adjusted using the two families of sextupoles which were placed on the quadrupoles.



## Summary

- The SCT can be optimized by varying the chromaticity of the ring. One point of **optimal chromaticity** for SCT can be found.
- The reason why the maximal SCT is not at  $\xi_x = 0$ ,  $\xi_y = 0$  is only partially understood and related to **intrinsic spin resonances**. Further investigation is needed.

## References

- [1] T. Fukuyama and A. J. Silenko, Derivation of Generalized Thomas-Bargmann-Michel-Telegdi Equation for a Particle with Electric Dipole Moment, Int. J. Mod. Phys A28, 1350147, 2013.
- [2] D. C. Sagan, Bmad: A relativistic charged particle simulation library, Nuclear Instruments and Methods in Physics Research A, vol.558, pp.356-359, 2006.
- [3] A. Lehrach, S. Martin and R. Talman, Design of a Prototype EDM Storage Ring, Proceedings of Science, PoS SPIN2018, 144, 2018.

Contact: Maximilian Vitz - m.vitz@fz-juelich.de